



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PUBLIC HEALTH REPORTS.

UNITED STATES.

Statistical reports of States and cities of the United States—Yearly and monthly.

CALIFORNIA—*Oakland*.—Month of October, 1902. Estimated population, 75,000. Total number of deaths, 90, including diphtheria, 7; enteric fever, 1, and 6 from phthisis pulmonalis.

CONNECTICUT—*Hartford*.—Month of October, 1902. Estimated population, 81,000. Total number of deaths, 92, including enteric fever, 2, and 8 from phthisis pulmonalis.

ILLINOIS—*Bloomington*.—Month of October, 1902. Estimated population, 25,000. Total number of deaths, 25, including enteric fever, 1, and 2 from tuberculosis.

Peoria.—Month of October, 1902. Census population, 56,100. Total number of deaths, 53, including enteric fever, 5, and 3 from tuberculosis.

LOUISIANA—*New Orleans*.—Month of October, 1902. Estimated population, 305,000—white, 223,500; colored, 81,500. Total number of deaths, 462—white, 289; colored, 173—including diphtheria, 3; enteric fever, 8; whooping cough, 2, and 91 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended November 8, 1902, from 85 observers, indicate that pleuritis, scarlet fever, remittent fever, and measles were more prevalent and diarrhea, enteric fever, and diphtheria were less prevalent than in the preceding week. Meningitis was reported present at 4, whooping cough at 20, smallpox at 20, measles at 21, diphtheria at 40, scarlet fever at 77, enteric fever at 31, and phthisis pulmonalis at 225 places.

MONTANA—*Helena*.—Month of October, 1902. Estimated population, 13,000. Total number of deaths not reported. One death from scarlet fever reported.

NEW HAMPSHIRE—*Manchester*.—Month of October, 1902. Census population, 56,987. Total number of deaths, 113, including diphtheria, 9; enteric fever, 1; measles, 3, and 2 from tuberculosis.

NEW JERSEY—*Passaic*.—Three weeks ended November 8, 1902. Census population, 27,777. Total number of deaths, 34, including 1 from diphtheria.

NEW YORK—*Saratoga*.—Month of October, 1902. Census population, 12,119. Total number of deaths, 17, including 2 from tuberculosis.

PENNSYLVANIA—*Dunmore*.—Month of October, 1902. Estimated population, 15,000. Total number of deaths, 17, including 1 from enteric fever.

TENNESSEE—*Nashville*.—Month of October, 1902. Estimated population, 81,320—white, 51,128; colored, 30,192. Total number of deaths, 136—white, 77; colored, 59—including diphtheria, 1; enteric fever, 6; scarlet fever, 1, and 20 from tuberculosis.

UTAH—*Ogden*.—Month of October, 1902. Estimated population, 20,000. Total number of deaths, 18. No deaths from contagious diseases.